




Synthesis, model validation, and data-assimilation on centennial time-scales

Michael Dietze, Jason McLachlan, Steve Jackson, Simon Goring,
Chris Paciorek, Jack Williams, and PaleEON team members


February 5, 2013


NACP 4th All Investigators Meeting, Albuquerque, NM

Biology drives Physics



Biology drives Physics







Jason McLachlan	Notre Dame
Michael Dietze	Boston University
Steve Jackson	U. Arizona
Chris Paciorek	UC Berkeley
Jack Williams	U. Wisconsin

60+ PaleEON team members

PaleON Goals

- Validation

- *How well do current models simulate decadal-to-centennial ecosystem dynamics when confronted with past climate change, and what factors most limit model accuracy?*

- Inference

- *What net carbon fluxes are compatible with an observed species composition and disturbance regime? Was the terrestrial biosphere a carbon sink or source during the Little Ice Age and Medieval Climate Anomaly?*


- Initialization

- *How sensitive are ecosystem models to initialization state and equilibrium assumptions? Do data-constrained simulations of centennial-scale forest dynamics improve 20th-century simulations?*

- Improvement

PaIEON Approach

Pre-PaIEON






PaIEON Approach


Pre-PaIEON

Current


PaIEON 2
(proposed)








(Goring, Mladenoff, Cogbill)






INTERIOR—GEOLOGICAL SURVEY, RESTON, VIRG. NIA—1980

Public Land Survey System (PLSS)


Township 2 South Range 3 West

Section 14




Section 14 shows both normal division of the section into aliquot parts and the fractional division into government lots.

Historical Vegetation




Goring in prep







Potential Vegetation




Ramankutty & Foley

Species Composition






STEPPS





An example from New England





Key features of past 2000 years

- Arrival of American chestnut
- Decline in beech and hemlock
- Spatial persistence of ecotone
- 19th century land-use impact




(Fuller et al 1998)


Statistical estimates of changing forest composition




Non-stationary trends in composition




Phase 1: Validation





Temperate Broadleaf Deciduous
850 AD
CMIP5/PMIP3 “Last Millennium”




C3 Grass
850 AD

Plant Functional Type Grid Fraction

Year 850; PFT 9 C3 Crop

C3 Crop
850 AD





Plant Functional Type Grid Fraction (%)

Equiangular (Regional) projection centered on -92.00°E 46.00°N

Data Min = 0.0E+00, Max = 8.7E+01

Centennial Sensitivity Analysis


ED2 model, Harvard Forest



Brett Raczka & Ken Davis, PSU
See Poster 128

Phase 2: Assimilation

State-Variable Data Assimilation



$$P(\theta|y) \propto P(y|\theta) P(\theta)$$

Updated State

Data

Model

UNDERC



UNDERC



Crampton Lake

Steigerwaldt Land Services



Pickerel Creek

Treehaven

5 -

Three 0.05ha plots

Data: Sam Pecoraro

6 - Prairie Peninsula


za Prairie Biological Station



University of Kansas Field Station



DBH


Increment





TREE GROWTH INFERENCE AND PREDICTION FROM DIAMETER CENSUSES AND RING WIDTHS



JAMES S. CLARK,^{1,2,3,4,6} MICHAEL WOLOSIN,^{2,3} MICHAEL DIETZE,^{2,3} INÉS IBÁÑEZ,^{2,3} SHANNON LADEAU,^{2,3,7}
 MIRANDA WELSH,¹ AND BRIAN KLOEPPEL⁵













NEE



GPP



Phase 2 Goals

- Proof of concept:
Assimilate 1000+ yrs for 5 sites
 - Data: pollen proxy, tree rings, settlement & modern inventory
 - Drivers: CMIP5 GCM downscaled realizations
 - Ensemble of CMIP5 GCMs
 - Ensemble of downscaled realizations
- Expand to regional scale
 - Inference
 - Initialization for the modern

Where we are now...

- GCM downscaling more complex and time consuming than anticipated...
- Very little proxy data independent of vegetation

PaIEON2

- New PLS
- Experimental design
- HIPS
- Tree rings
- Charcoal
- Paleo-climate

