

CLASS NUMBER THROUGH ADÈLES

TING GONG

CONTENTS

1. Introduction	1
2. \mathfrak{p} -adic Numbers	1
2.1. Norms and \mathfrak{p} -adic Integers	1
2.2. Completion of O_v	2
3. Adeles and Ideles	3
3.1. Properties of Adeles	3
3.2. Properties of Ideles	5
4. Applications in Algebraic Number Theory	6
4.1. The Idele Class Group	6
4.2. Approximation Theorem	7
4.3. Finiteness of Class Number	7
References	8

1. INTRODUCTION

In this paper, we are going to prove the Dirichlet's Unit Theorem and the finiteness of ideal class group through adèles and idèles. This introduces a more topological method to prove the results in algebraic number theory by using knowledge in local field theory and global field theory.

2. \mathfrak{p} -ADIC NUMBERS

2.1. Norms and \mathfrak{p} -adic Integers.

Definition 2.1. Let F be a field, $x, y \in F$, a map $\|\cdot\| : F \rightarrow \mathbb{R}_{\geq 0}$ satisfying the following properties is called a *norm*:

- (i) $\|x\| = 0$ if and only if $x = 0$
- (ii) $\|x \cdot y\| = \|x\| \cdot \|y\|$
- (iii) $\|x + y\| \leq \|x\| + \|y\|$

Definition 2.2. A norm is called *non-Archimedean* if $\|x+y\| \leq \max(\|x\|, \|y\|)$

Definition 2.3. Let F be a field, $x, y \in F$, a *valuation* is a map $v : F \rightarrow \mathbb{R}$ with the following properties:

Date: Started September 26, 2018. Last revision November 14, 2019.

- (i) $v(x) = \infty$ if and only if $x = 0$
- (ii) $v(xy) = v(x) + v(y)$
- (iii) $v(x + y) \geq \min(v(x), v(y))$

Definition 2.4. A valuation v is called *discrete* if the image of v is $\alpha\mathbb{Z}$ for some $\alpha \geq 0$

Definition 2.5. If k is a number field, O_k its ring of integers, \mathfrak{p} a prime ideal of O_k , We call the map $v_{\mathfrak{p}} : k^{\times} \rightarrow \mathbb{Z}$ the *\mathfrak{p} -adic evaluation* if p generates \mathfrak{p} , $a \in k$, $a = up^m$, where u is a unit. Then $v_{\mathfrak{p}}(a) = m$ or $v_{\mathfrak{p}}(a) = \infty$ when $a = 0$.

Definition 2.6. Under the same construction as Def. 2.5, let f be the degree of the residue field extension O_k/\mathfrak{p} over $\mathbb{Z}/(p)$, $\|\cdot\|_{v_{\mathfrak{p}}} : k \rightarrow \mathbb{R}$ is called the *\mathfrak{p} -adic norm*, if $\|a\|_{v_{\mathfrak{p}}} = \|\mathfrak{p}\|^{-fv_{\mathfrak{p}}(a)}$, or $\|a\|_{v_{\mathfrak{p}}} = 0$ when $a = 0$

We note that $\|\cdot\|_{v_{\mathfrak{p}}}$ is a non-archimedean norm.

Definition 2.7. We define $O_{\mathfrak{p}}$ to be the localization of O_k at \mathfrak{p} . Thus $O_{\mathfrak{p}} = \{\frac{a}{b} | b \notin \mathfrak{p}, a, b \in O_k\}$

Consider $M_n = O_{\mathfrak{p}}/(p^n)$, and the map $\varphi_m : M_n \rightarrow M_{n-1}$, by $\varphi(m_n) = m_n \bmod p^{n-1}$, and we see that $M_{n-1} \hookrightarrow M_n$ and that $\{M_n\}$ forms an projective system.

Definition 2.8. We define the inverse limit $\varprojlim M_n$ to be the *\mathfrak{p} -adic integers*, we denote as O_v .

Remark 2.9. $O_k \subseteq O_{\mathfrak{p}} \subseteq O_v$. This is easy to check, $O_k \subseteq O_{\mathfrak{p}}$ is clear since one can pick $b = 1$. Consider $a, b \in O_k$, then $a, b \in O_v$ by map $a \mapsto M_{v_{\mathfrak{p}}(a)}$ and similarly for b . Then one can do long division to see that $a/b \in O_v$. Thus $O_{\mathfrak{p}} \subseteq O_v$

Definition 2.10. We define A_k to be the *canonical set* of k , which is a set of all the absolute values, one non-archimedean absolute value for each \mathfrak{p} , and an archimedean absolute value for each real and complex embedding. We denote the set of archimedean absolute values as S_{∞} .

2.2. Completion of O_v . In this section and throughout the rest of the paper, we denote $v = v_{\mathfrak{p}}$.

Remark 2.11. We observe that the units of O_v are the elements of O_v with valuation 0, otherwise they all divide p , the uniformizing element. Therefore, we get the field of fraction of O_v by inverting p . We write it as $K_v = O_v(p^{-1})$

Definition 2.12. The *basic open sets* of K_v are defined as the open balls $B(x, r) = \{y \in K_v | \|x - y\|_v < r\}$

Proposition 2.13. *The basic open sets of K_v are closed, and K_v is totally disconnected.*

Proof. It suffices to check that $B(0, r)$ are closed. Consider the image of $\|a\|_v = \|\mathfrak{p}\|^{-v_{\mathfrak{p}}(a)}$, where $v_{\mathfrak{p}}$ has integer image. Thus $\text{im}(\|\cdot\|_v) = \{\|\mathfrak{p}\|^{\mathbb{Z}}\}$. Therefore, we see that the image is discrete besides at 0, which is a limit point. Thus, for some small $\varepsilon > 0$, $B(0, r) = \overline{B(0, r - \varepsilon)}$. Therefore, the basic open sets are closed.

Then we prove K_v is totally disconnected by showing if $a \in K_v$, the connected component C_a is the disjoint union of two open subsets. Let $a \in C_a$, but $C_a \neq \{a\}$. Thus $\exists n > 0$ such that $B(a, p^{-n}) \cap C_a \neq \emptyset$, but $C_a \neq B(a, p^{-n})$. Thus $C_a = (B(a, p^{-n}) \cap C_a) \sqcup ((K_v - B(a, p^{-n})) \cap C_a)$. Therefore, K_v is totally disconnected. \square

Theorem 2.14. *K_v is a locally compact, complete topological field with compact open and closed subsets O_v and O_v^**

Proof. It is clear that K_v is a topological field, considering term by term addition and multiplication to be continuous. Completeness is easy to check since one can check if convergent sequences are Cauchy. Since K_v is discrete, and every point is contained in a ball, which is finite. Thus, each basic open sets is compact, K_v is locally compact.

Since each M_n is discrete and finite, thus compact, by Tychonoff theorem, $\prod_{n=1}^{\infty} M_n$ is compact. Then it suffices to show that O_v is closed in $\prod_{n=1}^{\infty} M_n$ for O_v to be compact. If $(a_m) \notin O_v$, then $\exists N > 0$ such that $\varphi_N(a_N) \neq a_{N-1}$. Thus $(a_m) \in (a_1, \dots, a_{N-1}) \times \prod_{m>N} M_m$, which is open. Thus The complement of O_v is open, thus O_v is closed.

Since $O_v^* = O_v - \mathfrak{p}_v = O_v - B(0, 1)$, we see that O_v^* is closed, and O_v^* is a subspace of O_v , which is compact. Thus O_v^* is compact. \square

Proposition 2.15. *O_v^*, O_v, K_v are topological closures of O_k^*, O_k, k respectively.*

3. ADELES AND IDELES

3.1. Properties of Adeles.

Definition 3.1. We define *adeles* \mathbb{A}_k to be the subset of $\prod_{v \in A_k} k_v$, such that $\mathbb{A}_k = \{(a_v)_{v \in A_k} \mid a_v \in O_v \text{ for all but finitely many } v\}$

Remark 3.2. For some S containing S_{∞} and $S \subset A_k$, the basic open sets of \mathbb{A}_K is defined as $\prod_{v \in S} U_v \times \prod_{v \notin S} O_v$, where U_v is open in K_v . In this fashion we define the topology of \mathbb{A}_k

Proposition 3.3. *\mathbb{A}_k is a locally compact topological ring.*

Proof. We first show \mathbb{A}_k is locally compact. Since the basic open sets of \mathbb{A}_k are in the form $\prod_{v \in S} U_v \times \prod_{v \notin S} O_v$, we know that $\mathbb{A}_k \subset \prod_{v \in S} K_v \times \prod_{v \notin S} O_v$. Since each O_v is compact, $\prod_{v \notin S} O_v$ is compact by Tychonoff theorem, and each K_v is locally compact, thus $\prod_{v \in S} K_v$ is locally compact. Therefore, we know that $\prod_{v \in S} K_v \times \prod_{v \notin S} O_v$ is locally compact, thus \mathbb{A}_k is locally compact.

Next, \mathbb{A}_k is a ring is clear. We check continuity. Consider $f(a, b) = a + b$. Pick a basic open set $U = \prod_{v \in S} B(c_v, r_v) \times \prod_{v \notin S} O_v$. If $(a_v, b_v) \in f^{-1}(U)$. Then $\|a_v + b_v - c_v\|_v < r_v$. Next pick the sets $U_1, U_2 \subset A$ such that $a_v \in U_1, b_v \in U_2, f(U_1), f(U_2) \subset U$, then $U_1 = \prod_{v \in S} B(a_v, r_v/4) \times \prod_{v \notin S} O_v$ and $U_2 = \prod_{v \in S} B(b_v, r_v/4) \times \prod_{v \notin S} O_v$ works, since $f(U_1 \times U_2) = \prod_{v \in S} B(a_v + b_v, r_v/2) \times \prod_{v \notin S} O_v \subset \prod_{v \in S} B(c_v, r_v) \times \prod_{v \notin S} O_v$. Similarly, consider $f(a, b) = ab$, and the balls with radius $\sqrt{r_v/2}$. \square

Proposition 3.4. *Let $\iota : k \hookrightarrow \mathbb{A}_k$ such that $\iota(x) = x_v$, then $\iota(k)$ is a discrete subring of \mathbb{A}_K*

Proof. k is a subring is clear. We prove that it is discrete. Consider the basic open set at 0: $U = \prod_{v \in S_\infty} B(0, 1/2) \times \prod_{v \notin S_\infty} O_v$. Assume $0 \neq \alpha \in U \cap k$, then $\|\alpha\|_v \leq 1$ for non-archimedean evaluation and $\|\alpha\|_v < 1/2$ for archimedean ones. Thus $\prod_{v \in A_k} \|\alpha\|_v^{n_v} < 1/2$. But $\prod_{v \in A_k} \|\alpha\|_v^{n_v} = 1$ by product formula. Thus contradiction. Thus $U \cap k = 0$. Thus $\iota(k)$ is discrete. \square

Proposition 3.5. *Consider $\lambda : K_v \hookrightarrow \mathbb{A}_K$, if $x \in K_v$, then $\lambda(x) = (0, \dots, 0, x, 0, \dots) \in \mathbb{A}_K$. $\lambda(K_v)$ a closed subring, and it inherits the usual topology on K_v .*

Proof. The inheritance of the topology is clear. Now we prove that $\lambda(K_v)$ is closed. Let $a \in \mathbb{A}_K - \lambda(K_v)$, then $\exists v' \neq v$ such that $a_{v'} \neq 0$. Consider the following open set: $B(a_v, \|a_v\|/2) \times \prod_{w \in A_k - S_\infty, w \neq v'} O_w \times \prod_{v'' \in S_\infty} K_{v''}$. This is disjoint to $\lambda(K_v)$. Thus $\lambda(K_v)$ is closed. \square

Next, we are going to show that \mathbb{A}_K/k is compact.

Lemma 3.6. *$k + \mathbb{A}_K^{S_\infty} = \mathbb{A}_K$, where $\mathbb{A}_K^{S_\infty}$ is the set $\prod_{v \in S_\infty} K_v \times \prod_{v \notin S_\infty} O_v$. Thus we have any adele can be written as a sum of an element of k and $\mathbb{A}_K^{S_\infty}$.*

Proof. We prove if $a \in \mathbb{A}_K$, then $\exists \alpha \in k$ such that $\forall v \notin S_\infty, a_v - \alpha \in O_v$. Since $a_v \in O_v$ for all but finitely many $v \notin S_\infty$, we take c such that $v_p(c) > -v_p(a)$, where p is a prime of \mathbb{Z} lying under \mathfrak{p} with $a_v \notin O_{v_p}$, and such c exists since there are finitely many such p . Therefore, $ca \in O_v$.

Then let S be the set of primes of O_k dividing cO_k , by approximation theorem, for any distinct $v_1, \dots, v_s \in A_k, \alpha_1, \dots, \alpha_s \in k, \varepsilon > 0, \exists \alpha \in k$ such that $\|\alpha - \alpha_i\|_{v_i} \leq \varepsilon$. Therefore, let $\alpha \in O_k$ with $\alpha \equiv ca_v \pmod{\mathfrak{p}^m}$, for $\mathfrak{p} \in S$ and large m . Therefore, if $\mathfrak{p} \notin S$, then $c \in O_v^*$, so $\alpha/c \in O_v, a_v - \alpha/c \in O_v$. If $\mathfrak{p} \in S$, then if $m > v_p(c), a_v - \alpha/c \in O_v$. \square

Theorem 3.7. *\mathbb{A}_k/k is compact.*

Proof. Consider the restricted mapping $\iota|_{K_v} : k \hookrightarrow \prod_{v \in S_\infty} K_v = \mathbb{R}^n$. And O_v , as the \mathfrak{p} -adic integers, forms the lattice. Let $P \in \mathbb{R}^n$ be a fundamental parallelepiped for the lattice O_k , since it has the same rank as \mathbb{R}^n , P

is bounded and \bar{P} is compact. Also, we proved that O_v is compact. By Tychonoff theorem, $Q = \prod_{v \notin S_\infty} O_v \times \bar{P}$ is compact. By lemma 3.6, let $a \in \mathbb{A}_k$, pick $\alpha \in k$ such that $a - \alpha \in \mathbb{A}_K^{S_\infty}$, we translate $a - \alpha$ by β to get $a - \alpha + \beta \in \prod_{v \notin S_\infty} O_v \times \bar{P}$, and $-\alpha + \beta \in k$ which translates a into a compact set. Thus let $\pi : \mathbb{A}_k \rightarrow \mathbb{A}_k/k$, then $\pi(Q) = \mathbb{A}_k/k$ which is compact. \square

3.2. Properties of Ideles.

Definition 3.8. We define *ideles* \mathbb{I}_K to be the subset of $\prod_{v \in A_k} K_v^*$, such that $\mathbb{I}_K = \{(a_v)_{v \in A_k} | a_v \in O_v^* \text{ for all but finitely many } v\}$.

Remark 3.9. One can see that the ideles is the units in adeles. However, since the adeles forms a ring, the units of a ring doesn't need to be a multiplication group since the inverse doesn't need to be continuous. And there are two ways to define its basic open set.

- (1) One can define the topology with basic open sets similarly to that of the adeles. For some finite $S \subset A_k$, and $S_\infty \subset S$, the basic open sets of the ideles are $\prod_{v \in S} U_v \times \prod_{v \notin S} O_v^*$ where U_v is open in K_v^* for each $v \in S$.
- (2) One can also consider $\phi : \mathbb{I}_K \rightarrow \mathbb{A}_k \times \mathbb{A}_k$ with the subspace topology in $\mathbb{A}_k \times \mathbb{A}_k$, where $\phi(a) = (a, a^{-1})$. We observe that ϕ is a homeomorphism. Let $U = (\prod_{v \in S} U_v \times \prod_{v \notin S} O_v^*, \prod_{v \in S} U_v \times \prod_{v \notin S} O_v^*)$. Then $U \cap \phi(\mathbb{I}_K) = (\prod_{v \in S} U_v \cap (U'_v)^{-1} \times \prod_{v \notin S} O_v^*, \prod_{v \in S} (U_v^{-1} \cap U'_v) \times \prod_{v \notin S} O_v)$, whose preimage is $\prod_{v \in S} U_v \cap (U'_v)^{-1} \times \prod_{v \notin S} O_v^*$, which is open. Thus we have ϕ is homeomorphic.

Proposition 3.10. \mathbb{I}_K is a topological group.

Proof. Since \mathbb{I}_K is the kernel of our normed map, it is clearly a group.

Consider $\phi : \mathbb{I}_K \hookrightarrow \mathbb{A}_k \times \mathbb{A}_k$. Since we have shown ϕ is a homeomorphism, ϕ^{-1} is continuous, thus the inversion map is continuous.

Then define $\psi : \mathbb{A}_k \times \mathbb{A}_k \times \mathbb{A}_k \times \mathbb{A}_k \rightarrow \mathbb{A}_k \times \mathbb{A}_k$ by the following construction: let $(a_1, a_2), (a_3, a_4) \in \mathbb{I}_K \subset \mathbb{A}_k \times \mathbb{A}_k$, then $(a_1, a_2)(a_3, a_4) = (a_1 a_3, a_2 a_4)$. This function ψ is continuous since \mathbb{A}_k is an adelic ring, thus a topological ring. \square

Proposition 3.11. Let $\iota : k^* \hookrightarrow \mathbb{I}_K$ such that $\iota(x)_v = x \in k_v^*, \forall v$, then $\iota(k^*)$ is a discrete subgroup of \mathbb{I}_K

Proof. Consider the basic open set at 1: $U = \prod_{v \in S_\infty} B(1, 1/2) \times \prod_{v \notin S_\infty} O_v^*$. Assume $0 \neq \alpha \in U \cap k^*$, then $\|\alpha - 1\|_v \leq 1$ for non-archimedean evaluation and $\|\alpha - 1\|_v < 1/2$ for archimedean ones. Thus $\prod_{v \in A_k} \|\alpha - 1\|_v^{n_v} < 1/2$, which contradicts the product formula. Thus $U \cap k^* = 1$. Thus $\iota(k)$ is discrete. \square

Proposition 3.12. Consider $\lambda : K_v^* \hookrightarrow \mathbb{I}_K$, if $x \in K_v^*$, then $\lambda(x) = (1, \dots, 1, x, 1, \dots) \in \mathbb{I}_K$. $\lambda(K_v^*)$ a closed subgroup.

Proof. Let $a \in \mathbb{I}_k - k_v^*$ and choose $v' \neq v$, $a_{v'} \neq 0$. Consider $B(a_v, \|a_v - 1\|/2) \times \prod_{A_k - S_\infty, n \neq v'} O_n^* \times \prod_{n \in S_\infty} K_n^*$, which is an open neighborhood disjoint from $\lambda(K_v^*)$. Thus $\lambda(K_v^*)$ is closed. \square

4. APPLICATIONS IN ALGEBRAIC NUMBER THEORY

4.1. The Idele Class Group.

Remark 4.1. We define a map: $\|\cdot\| : \mathbb{I}_K \rightarrow \mathbb{R}^+$ with $\|a\| = \prod_{v \in A_k} \|a_v\|_v^{n_v}$

Lemma 4.2. $\|\cdot\|$ is continuous.

Proof. Assume $(b_0, b_1) \subset \mathbb{R}^+$. We show that the preimage is open in \mathbb{I}_K . If $a \in \mathbb{I}_K$, and $\|a\| \in (b_0, b_1)$, then pick an archimedean valuation v_0 . Let S be the set of valuations where a is not in O_v^* , consider the open sets $U_r = \prod_{v \in S, v \neq v_0} B(a_v, 1) \times \prod_{v \notin S} O_v^* \times B(a_{v_0}, r)$ where r varies. Thus $a \in U_r$. Then we can make r small so that $\|U_r\|$ lies in (b_0, b_1) . \square

Remark 4.3. Since $\{1\}$ is closed in \mathbb{R}^+ , $\ker(\|\cdot\|)$ is a closed subgroup in \mathbb{I}_K . We call this \mathbb{I}_K^1 . By product formula and Proposition 3.12, $k^* \subseteq \mathbb{I}_K^1$, is a discrete subgroup.

Now, let \mathfrak{F}_K be the multiplicative group of fractional ideals of k . Let $\mathfrak{P}\mathfrak{F}_K$ be the subgroup of principal ideals. Let $C_K = \mathfrak{F}_K / \mathfrak{P}\mathfrak{F}_K$ be the class group over k .

Theorem 4.4. $\mathbb{I}_K / k^* \mathbb{I}_K^{S_\infty} \cong C_K$

Proof. We define $\phi : \mathbb{I}_K \rightarrow \mathfrak{F}_K$ by $\phi(a) = \prod_{v \in A_k - S_\infty} \mathfrak{p}^{v_p(a_v)}$. Since only finitely many $v_p(a_v)$ are non-zero, this is a fractional ideal. Since \mathfrak{F}_K is generated by \mathfrak{p} , where \mathfrak{p} varies over primes of O_k . Define $a_{\mathfrak{p}} \in \mathbb{I}_k$ with $(a_{\mathfrak{p}})_{v_{\mathfrak{p}}} = p_v$, the uniformizer for $k_{v_{\mathfrak{p}}}$ and $(a_{\mathfrak{p}})_{v'} = 1$ if $v' \neq v$; thus $\phi(a_{\mathfrak{p}}) = \mathfrak{p}$. Therefore, the map ϕ is surjective. The kernel of this map happens when all $v_p(a_v) = 0$, but the archimedean entries can vary. Thus the kernel is $\mathbb{I}_K^{S_\infty}$. Let $\alpha \in k^*$, $(\alpha) = \alpha O_k$, then $\phi(\alpha)$ is (α) . Thus $\phi(k^*) \subset \mathfrak{P}\mathfrak{F}_K$. Thus $\phi(k^*) = \mathfrak{P}\mathfrak{F}_K$. Thus, there is an induced surjective homomorphism $\psi : \mathbb{I}_K / k^* \rightarrow C_K$. Consider $\varphi : \mathbb{I}_K \rightarrow C_K$, $\varphi(a) = \phi(a) \mathfrak{P}\mathfrak{F}_k$, then φ is surjective since ϕ is surjective. Let $a \in \ker(\varphi)$, then $\phi(a) \in \mathfrak{P}\mathfrak{F}_K$. Thus $\phi(a) = \alpha O_k$ for some $\alpha \in k^*$. And since $\alpha O_k = \phi(\iota(\alpha))$, so $\phi(a) = \phi(\iota(\alpha))$. Thus $\phi(a \iota(\alpha)^{-1}) = 1$. Thus, $a \iota(\alpha) \in \mathbb{I}_K^{S_\infty}$. Thus $a \in \iota(\alpha) \mathbb{I}_K^{S_\infty}$. Thus $\ker(\varphi) \subset k^* \mathbb{I}_K^{S_\infty}$. The other inclusion is clear by the discussion above. Then $\ker(\varphi) = k^* \mathbb{I}_K^{S_\infty}$. Thus by the first isomorphism theorem, $\mathbb{I}_K / k^* \mathbb{I}_K^{S_\infty} \cong C_K$ \square

Definition 4.5. We call $C = \mathbb{I}_K / k^*$ the idele class group, and $C^1 = \mathbb{I}_K^1 / k^*$.

Definition 4.6. Let S be a finite subset of A_k containing S_∞ , we call $k_S = \mathbb{I}_K^S \cap k^*$ the S -units of k . And we call \mathbb{I}_K^S / k_S the group of S -idele classes and denote by C_S . We set $\mathbb{I}_K^{S^1} = \mathbb{I}_K^S \cap \mathbb{I}_K^1$ and $C_S^1 = \mathbb{I}_K^{S^1} / k_S$

Remark 4.7. For each S , $C_S \hookrightarrow C$. Since \mathbb{I}_K^S is both open and closed in \mathbb{I}_K , C_S is both open and closed in C . Similarly, $C_S^1 \hookrightarrow C$, and C_S^1 is both open and closed in C^1 . Then consider the map $C \rightarrow C_K$, then the kernel will be C_{S_∞} since $\mathbb{I}_K/k^*\mathbb{I}_K^{S_\infty} \cong C_K$, by third isomorphism theorem, $\mathbb{I}_K/k^*\mathbb{I}_K^{S_\infty} \cong \frac{\mathbb{I}_K}{k^*}/\frac{\mathbb{I}_K^{S_\infty}k^*}{k^*}$, and the latter is isomorphic to C_{S_∞} by the second isomorphism theorem. Thus $C/C_{S_\infty} \cong C_K$

Finally, notice the map $C^1 \rightarrow C_K$ is also surjective, since if $\mathfrak{f} \in \mathfrak{F}$, $a \in \mathbb{I}_K$ with $\phi(a) = I$, we can change an archimedean value to get $a' \in \mathbb{I}_K^1$, and $\phi(a') = I$. Thus the kernel is $C_{S_\infty}^1$. So we have $C^1/C_{S_\infty}^1 \cong C_K$

4.2. Approximation Theorem.

Definition 4.8. Let $a \in \mathbb{A}_K$, we define $L(a) \subseteq k$, such that $L(a) = \{\alpha \mid \|\alpha\|_v \leq \|a\|_v, \alpha \in k, \forall v \in A_k\}$. We write $\lambda = |L(a)|$.

Remark 4.9. If $\alpha \in k^*$, $x \in L(a)$, then since $\|x\|_v \leq \|x\|_v$, then $\|\alpha x\|_v = \|\alpha\|_v \|x\|_v \leq \|\alpha\|_v \|x\|_v = \|\alpha x\|_v$. Therefore, there is a bijection between $L(a)$ and $L(\alpha a)$, thus $\lambda(a) = \lambda(\alpha a)$

Theorem 4.10. $\exists c_0$ a constant, depending only on k such that for any $a \in \mathbb{A}_K$, $\lambda(a) \geq c_0 \|a\|$.

Proof. For the proof of this theorem, refer to [1]Theorem 8.1. \square

Lemma 4.11. Let $a \in \mathbb{I}_K$, $\|a\| \geq 2/c_0$, then $\exists \alpha \in k^*$ such that we have $\forall v \in A_k, 1 \leq \|\alpha a\|_v \leq \|a\|$

Proof. For the proof of this lemma, refer to [1]Lemma 9.1 \square

4.3. Finiteness of Class Number.

Theorem 4.12. C^1 is compact

Proof. Let $\psi : \mathbb{I}_K \rightarrow \mathbb{R}^+$ by $\psi(a) = \|a\|$. Since if $a \in k^*$, $\phi(a) = 1$ by the product formula. Thus consider $\psi : C \rightarrow \mathbb{R}^+$. Thus $\ker(\psi) = C^1$. Then since if $\rho \in \mathbb{R}^+$, $a_\rho \in \mathbb{I}_K$ with $\|a_\rho\| = \rho$, then $\psi^{-1}(\rho) = a_\rho C^1$. Therefore, we see that ψ^{-1} is homeomorphic to C^1 . Thus we prove ψ^{-1} is compact.

Let $\rho > 2/c_0$, pick $a \in \psi^{-1}(\rho)$, then by Lemma 4.11, $\exists \alpha_a \in k^*$ such that $\forall v \in A_k, 1 \leq \|\alpha_a a\|_v \leq \rho$. Since $\|\cdot\|_v$ cannot take values between 1 and $N(\mathfrak{p})$, and there are only finitely many \mathfrak{p} with $N(\mathfrak{p}) \leq \rho$. Thus we have $\|\alpha_a a\|_{v_{\mathfrak{p}}} = 1$ for all but finitely many of $v_{\mathfrak{p}}$. Thus $\exists S \supset S_\infty$ such that $1 \leq \|\alpha_a a\|_v \leq \rho$ if $v \in S$ but $\|\alpha_a a\|_v = 1$ if $v \notin S$.

Now define $T = \prod_{v \in S} (\overline{B(0, \rho)} - B(0, 1)) \times \prod_{v \notin S} O_v^*$. Thus by Tychonoff theorem, T is compact, since $\psi^{-1}(\rho)$ is a close subset of $\phi(T)$, where $\phi : \mathbb{I}_K \rightarrow C$. Thus ψ^{-1} is compact. \square

Corollary 4.13. C_S^1 is compact for any finite set S containing S_∞ .

Theorem 4.14 (Finiteness of class number). For any number field k , C_K is finite.

Proof. Since $C^1/C_{S_\infty}^1 \cong C_K$, and C^1 is compact, thus C_K is compact. Since $C_{S_\infty}^1$ is open and for any topological group G , if H is an open subgroup of G , then G/H is discrete, then C_K is also discrete. Thus C_K is both compact and discrete, thus it is finite. \square

Lemma 4.15. *Any discrete subgroup Λ of \mathbb{R}^s is free abelian of rank $\dim \mathbb{R}\Lambda$*

Proof. We induct on $\dim \mathbb{R}\Lambda$. If $\dim \mathbb{R}\Lambda = 1$, since Λ is discrete, $\exists \lambda \in \Lambda$ closest to 0. Thus $\Lambda = \mathbb{Z}\lambda$.

Assume $\dim \Lambda = m$, let $\lambda_1, \dots, \lambda_m$ be a \mathbb{R} -basis for $\mathbb{R}\Lambda$. If Λ_0 is a subgroup of Λ spanned by $\lambda, \dots, \lambda_{m-1}$, then by induction hypothesis, $\Lambda_0 = \mathbb{Z}\lambda_1 \oplus \dots \oplus \mathbb{Z}\lambda_{m-1}$. Let $\lambda' \in \Lambda$ such that $\lambda = a_1\lambda_1 + \dots + a_m\lambda_m$ with $0 \leq a_i < 1$ for $i \leq m-1$ and $0 \leq a_m \leq 1$. Then λ' is bounded, thus finite. Let $\lambda' \in \Lambda'$ has minimal nonzero coefficient, assume $\lambda' = a'_1\lambda_1 + \dots + a'_m\lambda_m$. Assume if $\lambda \in \Lambda$, then $\exists t$ such that the m th coefficient of $\lambda - t\lambda'$ gives $0 \leq a_m < a'_m$. Then we modify by $\lambda_0 \in \Lambda_0$ to get $\lambda - t\lambda' - \lambda_0 \in \Lambda'$. Thus $a_m = 0$ since a'_m is minimal. Thus $\lambda - t\lambda' - \lambda_0 = 0$. Since $\lambda_1, \dots, \lambda_m$ are linearly independent, we have $\Lambda = \mathbb{Z}\lambda \oplus \dots \oplus \mathbb{Z}\lambda_m$. \square

Theorem 4.16 (Dirichlet Unit Theorem). *For any set finite set $S \in A_k$ of size s containing S_∞ , the S -units k_S have rank $s-1$.*

Proof. Let $v_1, \dots, v_s \in S$, and assume v_s is archimedean. Define $\log : \mathbb{I}_K^S \rightarrow \mathbb{R}^s$ by $\log(a) = (\log \|a_{v_1}\|_{v_1}^{n_{v_1}}, \dots, \log \|a_{v_s}\|_{v_s}^{n_{v_s}})$. Since it is continuous in each coordinate, \log is continuous. Also, since $a \in \mathbb{I}_K^{S_1}$, then $\|a\| = 1$, since $\|a_v\|_v = 1$ for $v \notin S$, $\log(\mathbb{I}_K^{S_1}) = \{x_1, \dots, x_s \in \mathbb{R} \mid x_1 + \dots + x_s = 0\} = H$. Thus $\dim \log(\mathbb{I}_K^{S_1}) = s-1$.

Since in a bounded region in \mathbb{R}^s , $\log(k_S)$ has bounded archimedean absolute values. Thus the coefficients of the polynomials of these elements over \mathbb{Z} is bounded. And the degree is bounded by $[k : \mathbb{Q}]$, there are finitely many such polynomials. Thus there are finitely many k maps into this bounded region. Thus $\log(k_S)$ is discrete. By the last lemma, $\log(k_S)$ is a free abelian group.

Let W be the subspace of H generated by $\log(k_S)$, then consider $\log : \mathbb{I}_K^{S_1}/k_S = C_S^1 \rightarrow H/W$. Since $\mathbb{I}_K^{S_1}$ generates H , the image generates H/W as an \mathbb{R} -vector space. Since \log is continuous, C_S^1 is compact, the image is compact. Then if H/W is non-trivial, then it has no non-trivial compact subgroups. Thus $H/W = 0, H = W, \dim \log(k_S) = s-1$. \square

Corollary 4.17. *The group of global units of a number field k is isomorphic to $\mu(k) \times \mathbb{Z}^{r_1+r_2-1}$, where $\mu(k)$ is the roots of unity, r_1 is the number of real embeddings, r_2 is the number of complex embeddings.*

REFERENCES

- [1] Weston, *The Idelic Approach to Number Theory*, 1999, Online access

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF NOTRE DAME, NOTRE DAME, IN
46556-4618.

Email address: `tgong@nd.edu`